

The resumption of the Spanish football league after COVID-19 breakdown: The model (on-going research: version 1)

Javier M. Buldú^{1,2}, Daniel Ruiz Antequera^{1,2} and Jacobo Aguirre³

¹ Complex Systems Group, Universidad Rey Juan Carlos, Camino del Molino s/n, 28943 Fuenlabrada, Madrid, Spain.

² Laboratory of Biological Networks, Centre for Biomedical Technology, UPM-Campus de Montegancedo, 28223 Pozuelo de Alarcón, Madrid, Spain.

³ Centro de Astrobiología (CSIC-INTA), Ctra. de Ajalvir km 4, 28850, Torrejón de Ardoz, Madrid, Spain.

The SEIR model applied to sport dynamics

In this work we make use of a discrete *SEIR Susceptible-Exposed-Infectious-Recovered* model [1] and adapt it conveniently to describe the propagation of COVID-19 during a football tournament. Specifically, we are concerned about the re-start of the Spanish national football league (*“La Liga”*), which is currently stopped with 11 fixtures remaining. The goal of the model is to simulate how COVID-19 would propagate in case the season was re-started after quarantine.

In SEIR models, a disease propagates through a network of individuals whose dynamical state can be either susceptible (healthy and susceptible to be infected), exposed (infected but in the incubation period and therefore unable to infect other individuals), infectious (infected and able to infect other individuals), and removed (which includes (i) recovered individuals after having suffered the infection and therefore immune and (ii) deceased people).




Figure 1. Schematic representation of the SEIR model adapted to a football competition. Players can be in different states: Susceptible (S), Exposed (E), Infectious (I) and Quarantined (Q). In case players are detected to be infected by the virus, they remain confined (indicated by the C suffix). After confinement, players will undergo a quarantine before being eligible to play again. Parameters β_{train} , β_{match} and β_{ext} account for the probability of being Exposed (E) during training, matches or externally (player social circle) respectively. Probability σ describes the transition from Exposed to Infectious (I). Probability γ controls the transition from Infectious to Recovered (R) or Quarantined (Q). Finally, γ_Q is related to the quarantine period a player must follow after recovery.

Figure 1 represents a sketch of the model. The individuals (players from now on) can be infected at any time from people different to the players (technical staff of the team, family, etc.) with a probability β_{ext} . The second source of infection occurs during the training period, where they can be infected from other players of their own team with probability β_{train} . Finally, during the matches, players are exposed to infection from the players of their own team and the adversary team with probability β_{match} .

Once a player has been infected and becomes exposed, he/she has a probability σ of finishing the incubation period and become infectious. Exposed and infectious players have, respectively, probabilities μ_E and μ_I of being detected as infected by COVID-19 via a virus test or because they show disease symptoms. If this is the case, players will be confined at their homes remaining in two possible states: exposed E_C or infectious I_C . Asymptomatic infectious players (belonging to class I, but not detected by virus tests), and confined infectious players, overcome the disease with probability γ . Note that confined players that have been recovered will remain quarantined (class Q) during a convalescence until they are prepared for playing again and become recovered (R) with probability γ_Q .

The frequency of virus tests N_{test}^{-1} and the frequency of matches N_{match}^{-1} are two critical variables for controlling the number of infected players during the championship, and therefore their influence in the model should be studied carefully. Note that the virus tests should be done in this context via Polymerase Chain Reaction (PCR) controls. The reason is that fast antibody or antigen detectors are only reliable more than a week after the infection, and in many cases even after the patient has already shown symptoms. This fact would allow the infectious (but not detected) players to spread the virus during several days, making the control of the disease a hard task.

Modeling the Spanish national league

Our model can be applied to a diversity of competitions related to team contact sports, but we have focused on the re-start of the Spanish male national league. Therefore, we considered a competition with $M = 20$ teams composed of $L = 25$ players, the latter being the upper limit of players that can be registered by a team in the competition. The generalization to *Liga Iberdrola* (Spanish feminine first division football league, with 16 teams), to the masculine or feminine football leagues of other countries, or even to other team sports (such as basketball, handball, rugby, etc.) is direct. Every team plays a match every N_{match} days, and during the $N_{match} - 1$ days in between the players train at their own stadiums. We supposed no resting days, as there is a clear interest for finishing the leagues as soon as possible, but including them in the model is trivial. We represented the training dynamics of the players, and contacts between them, using social networks instead of mean field contacts. In this way, players' social networks followed a random structure of connections and were generated using an Erdős-Rényi model [2] with a probability $p = 0.2$ of connecting two players. This was done to describe the internal professional and friendship dynamics that every player has during training times and also during lunch time, etc. During training time, the infectious players (class I) might infect their neighbours in the social network with probability β_{train} . During the match day, every infected player on the pitch can infect any other player of its own team or the adversary with probability β_{match} (here we used a mean-field approach due to the inevitable contact dynamics that players follow during a match). Note that players cannot avoid voluntarily the contact with other players in the contest (with the exception, perhaps, of celebrating a goal, that could be forbidden if necessary), and therefore the contagion probability during a match might be larger than expected at first glance.

There is a wide range of values in the recent literature regarding each of the parameters that define the different steps of the disease (see Table 1 for a summary of the parameters of the model). The latent period σ^{-1} is the average time from infection to infectiousness, the incubation period is the average time from infection to the appearance of the first symptoms, and the infectious period γ is the average time that the patient is infectious. Depending on the virus, the latent period can be shorter or larger than the incubation period. In the case of COVID-19 the latent period is 1 or 2 days shorter in average than the incubation period, which makes it specially easy for the disease to spread among the population during

Parameter	Value (days ⁻¹)
β_{exp}	1/100000
β_{train}	1/10
β_{match}	1/100
σ	1/3; Refs. [3, 4]
γ	1/5; Refs. [3, 4]
γ_Q	1/5
μ_E	$\mu_I/3$
μ_I	[0,1], 0.9

Table 1. Summary of the main parameters of the model.

the time in which people are infectious but asymptomatic. Regarding the mean incubation period, in [5] it was shown to be around 5 days, similar to that of SARS, and in [6] it was affirmed that it could be as short as 4 days. Note, however, that this quantity was not used in our model.

In [3] it was used a mean latent period σ^{-1} of 3 days and a mean infection period of $\gamma^{-1} = 5$ days, based on the Wuhan data. We selected these values because they were also used in other more recent studies [4]. Note, however, that these are mean values: in [7] it was shown that the probability that patients with mild symptoms infected other people was very low after a week from the appearance of symptoms, but these means that in mild cases of COVID-19 patients can be infectious for as much as 10 days.

The values of the probability μ_I of being detected as infectious, either because a player shows disease symptoms or because the virus test yields a positive result, has been considered to be within the window [0,1], being 0 in case of not doing any test and being asymptomatic and 1 when tests have 100% sensitivity. However, when sensitivity of the test is not analysed, we considered a value of 0.9 which is the typical one attributed to PCR tests. Concerning the probability of detecting an exposed individual, we set it as $\mu_E = \mu_I/3$, i.e. three times less than detecting an infected individual through the same test. The reason is that the viral load of an exposed individual is lower than that of an infectious one, therefore reducing the probability of a positive test result.

Furthermore, we have fixed the quarantine period σ_Q^{-1} to be 5 days, but varying slightly this quantity would not affect the model substantially. Also, there is no clear experimental data to fix the infecting probabilities β_{exp} , β_{train} and β_{match} . The probability β_{ext} of being infected from people different to the players will slowly decrease as more and more people in the country recover from the disease, but for simplicity we have supposed it constant during the whole league. The probabilities of being infected during a training day or a match have been fixed at moderate values and we checked that slightly varying them did not qualitatively change the results.

Finally, we simulated 1000 seasons using our discrete model and obtained the mean values and the standard deviations of each variable. Importantly, the seed of all simulations contained one player of the league who is already infected at the first day of the tournament. By doing so, the epidemic spreading begins at day one instead of any random day of the season.

Figure 2 shows one example of the results obtained with the model. In this case, we show the evolution of the number of players that have been infected when matches are played with a separation of 7 days. The three cases plotted in Fig. 2 correspond to: (i) not considering any test during the whole season, (ii) doing a PCR test (with 90% sensitivity every 7 days and (iii) doing a PCR test every day.




Figure 2. Number of players that have been infected at least one time at the end of the season. Lines correspond to: (red) not considering any test during the whole season, (blue) doing a PCR test (with 90% sensitivity every 7 days) and (green) doing a PCR test every day. Season contains 11 fixtures, i.e. just the ones remaining. Parameters of the simulation are indicated in Table 1, with $\mu_I = 0.9$. The seed of all simulations contained one player infected at the first day of the tournament.

References

1. Newman, M. *Networks: An Introduction* (Oxford University Press, Inc., New York, NY, USA, 2010).
2. Erdős, P., Rényi, A. *et al.* On random graphs. *Publicationes mathematicae* **6**, 290–297 (1959).
3. Wu, J. T., Leung, K. & Leung, G. M. Nowcasting and forecasting the potential domestic and international spread of the 2019-ncov outbreak originating in wuhan, china: a modelling study. *The Lancet* **395**, 689 – 697 (2020).
4. Lin, Q. *et al.* A conceptual model for the coronavirus disease 2019 (covid-19) outbreak in wuhan, china with individual reaction and governmental action. *International Journal of Infectious Diseases* **93**, 201–216 (2020).
5. Lauer, S. A. *et al.* The Incubation Period of Coronavirus Disease 2019 (COVID-19) From Publicly Reported Confirmed Cases: Estimation and Application (2020, date (06/03/2020)).
6. Guan, W.-j. *et al.* Clinical characteristics of 2019 novel coronavirus infection in china. *medRxiv* (2020). URL <https://www.medrxiv.org/content/early/2020/02/09/2020.02.06.20020974>. <https://www.medrxiv.org/content/early/2020/02/09/2020.02.06.20020974.full.pdf>.
7. Wolfel, R. *et al.* Virological assessment of hospitalized patients with covid-2019. *Nature* (2020).